

## Guidelines for avoiding antihistamines prior to allergy skin testing

Antihistamines must be discontinued prior to skin testing. Some antihistamines should be discontinued 48 hours prior to testing and some (as listed) need to be discontinued up to a week prior to testing. Decongestants without antihistamines can be taken up to the time of testing.

Please see the following list for some of the most common medications and when they should be discontinued. If you have any questions about antihistamines in your medication, please check with your physician or your pharmacist.

## TRICYCLIC ANTIDEPRESSANTS (STOP THESE 1 WEEK BEFORE)

Amitriptyline (Endep, Enovil, Elavil, Emitrip)	Limbitrol	
Amoxapine (Asendin)	Nortriptylline (Pamelor, Aventl)	
Desipramine (Norpramin, Pertofrane)	Protriptyline (Vivactil)	
Doxepin (Adapin, Sinequan)	Seroquel	
Imipramine (Janimine, Tipramine, Tofranil)	Trimipgramine (Surmontil)	

## **ANTIHISTAMINES** (STOP 48 HOURS – 1 WEEK BEFORE)

Actidil	Contac	Palgic – 1 week
Actifed	Coricidin	Pantanase
Alavert – 1 week	Cyproheptadine	Periactin – 1 week
Allegra – 1 week	Deconade	Phenergan – 5 days
Allegra D – 1 week	Deconamine	Polaramine
Allerest	Dexbrompheniramine	Polyhistinen D
Allergesic	Dexchlorpheniramine – 1 week	Promethazine
AllerX	Dimetane	Rondec
Antivert	Dimetapp	Rynatan
Astelin Nasal Spray	Dymista	Ryna – 12-S
Astepro	Diphenhydramine	Semprex
Atarax – 1 week	Doxylamine	Sinutab
Atrohist	Dramamine	Sudafed plus
Azelastine Nasal	Drixoral	Tanafed
Benadryl	Dristan	Tavist
Benylin	Extendryl	Trazodone
Bromfed	Fedahist	Triaminic
Brompheniramine	Fexofenadine – 1 week	TussiCaps – 1 week
Carbinoxamine maleate – 1 week	Flexril (muscle relaxer)	Tussionex pennkinetic – 1 week
Cetirizine	Formula 44	Vicks nyquil
Chlor-trimeton	Histex – 1 week	Vistaril
Chlorpheniramine	Hydroxyzine – 1 week	Xyzal – 1 week
Claritin – 1 week	Isoclor	Zyrtec
Claritin D – 1 week	Loratadine – 1 week	Zyrtec D
Clarinex – 1 week	Naldecon	ALSO, over the counter medication
		that contain antihistamines.
Clemastine fumarate	Olopatadine Nasal	
Clistin rondec	Optimine	
Clorpheniramine – 1 week	Orahist	
Comtrex	Ornade	